Introduction

For the last few decades, aesthetic dermatology has become increasingly popular. Most of our patients want to enhance their look without downtime. As aesthetic surgery continues to have a tremendous role in the growth of our specialty, cosmetic dermatology is growing, with many different techniques for “rejuvenation”. Among them, injection techniques with filler agents are a common part of everyday medical practice.

This special issue on fillers, written by a panel of international physicians, will help to guide everyone in this “jungle” and allow us to better evaluate risks and benefits of all these agents.

Available agents should be separated into two groups. The first one contains biodegradable products (Bovine collagen has been for a long time the gold standard), the longevity being variable but transitory. The second group contains non-biodegradable products (350 centistoke medical grade fluid silicone was utilized for a long time) whose longevity is supposed to be permanent.

For cosmetic indications (wrinkles treatment), it seems reasonable to utilize less risky products, which many feel are, for the time being, biodegradable. With the biodegradable products, in cases of complications, they will most often disappear spontaneously. With the permanent products, we have seen very severe reactions, which sometimes do not appear until a few years after the injection. Although new techniques (in particular with liquid injectable silicone) may help to obviate some of these long term complications, the

risk of permanent fillers will still be that of permanent complications.

The choice of the filler agent must be discussed with the patient according to the indications and the risks, which need to be exposed before any treatment is begun. Outside of the USA, regulations for launching a new medical device on the market do not require clinical studies with long term follow up. For this reason, in Europe we have seen a lot of severe complications with many filler agents (some of them are no longer on the market). In cases of severe complications, appearing sometimes several years after the injections, patients do not always remember which agent has been injected. In these cases, skin biopsy can prove by the histological aspect, the culpability of one of these filler agents.

The ideal filler has not yet been discovered (what is the ideal filler?), but over the last decade several new filler products have been developed and utilized. In addition, lipofilling continues to grow in popularity since there is no hypersensitivity, and results can be superb with proper technique.

Many of these new products and techniques, with the histological evidence and complications, will be discussed in this issue. We hope you find this international collaboration as enjoyable to read as we did to work on.

Pierre Andre, MD
Michael Kaminer, MD
Guest Editors