Introduction

In 1982, when the Food and Drug Administration approved injectable collagen for soft-tissue augmentation, the era of esthetic dermatology was born. In the decades that ensued, updates in esthetic dermatology largely consisted of new methods to cross-link bovine collagen or different methods of injecting it into various locations. When the Carruthers first reported the use of botulinum toxin for the treatment of dynamic rhytids in 1992, the next era of esthetic dermatology was begun. The use of botulinum toxins ignited interest within the dermatology community to explore the possibilities for nonsurgical esthetic improvement. This interest coincided with new techniques for performing liposuction that made the procedure safer and more effective, lasers that for the first time targeted specific components of the epidermis and dermis, injectable soft-tissue augmentation products that have increased persistence, cosmeceuticals based on a molecular understanding of aging skin and a host of other products, procedures and, perhaps most importantly, attitudes among physicians with respect to what can and cannot be done to enhance and improve appearance.

As with every paradigm shift, the use of botulinum toxins was greeted with skepticism in many parts of the dermatology community. It did not seem intuitively obvious to use a toxin to make people look better. However, as the publications demonstrating safety and efficacy increased, more physicians adopted Botox® into their clinical practices. At first, techniques were discussed to treat the glabella rhytids, and then periorbital wrinkles were treated. As dermatologists became more familiar with the intricacies of facial anatomy, the mid-face and lower face became amenable to treatment. Once the chin was being treated, the neck was not far behind. Presently, areas treated with botulinum toxins for cosmetic usage have increased to the point where most regions of the face are treated by some physicians. Newer toxins that use other formulations of type A toxin as well as type B toxins are being used in addition to the original type A formulation (Botox® Cosmetic, Allergan, Irvine, CA). Indications and technical considerations for botulinum toxin injections will be discussed in the article by Timothy Corcoran Flynn.¹

Until quite recently, products used for soft-tissue augmentation traditionally have included autologous fat (used for decades), collagens, and silicone. Newer technologies allowing for the cross linkage of hyaluronic acid molecules intro-

duced soft-tissue augmentation products that were safe and effective with a degree of persistence that allowed a long-term correction to be possible. Different iterations of the hyalurons became available in the United States in 2004, and many more are available in Europe and Canada. These products not only created a revolution in cosmetic dermatology in their own right but also have a synergistic role with the use of botulinum toxins. Combinations of the hyalurons and botulinum toxins are now the most common combination esthetic enhancement performed by most physicians. An update on hyaluronic acids is presented in the article by Mary Lupo.²

The increase in soft-tissue augmentation products is not limited to the hyalurons; newer materials include calcium hydroxylapatite and poly-L-lactic acids (PLAs). Presently approved for treatment of craniofacial defects and laryngeal injections, calcium hydroxylapatite (Radiesse™, San Mateo, CA) is expected to be approved for cosmetic soft-tissue augmentation. It was used by a variety of physicians in 2002 with a variety of outcomes. More recently, some consensus is developing regarding its utility for indications including nasolabial creases and marionette lines. Experience also has taught us where the product should not be injected, and both of these issues will be discussed in the article by Berlin et al.³

PLA (Sculptra; Dermik Laboratories) is presently approved for the treatment of HIV-associated lipoatrophy, but its popularity for esthetic indications is growing and it is likely that this indication will be obtained in the near future. Unlike other products, PLA is not a direct augmentation enhancer, that is to say, injecting 1 mL of PLA does not guarantee 1 mL of correction. Rather, the molecule stimulates collagen production and this replaces volume and can reposition the face. This product is a novel compound in its mechanism of action and potential for long-term autologous volume restoration. The uses of PLA will be reviewed in the article by Beer and Rendon.⁴

Esthetic dermatology also has been revolutionized by lasers and light sources. These devices, developed by dermatologists, first gained popularity for the treatment of vascular lesions. The use of alexandrite lasers for hair reduction increased the popularity of the devices within the dermatology community who, for the first time, had a safe and effective treatment for unwanted hair. One unfortunate offshoot of

this development was the adoption of this technology by those who lack the training or experience to perform these procedures and their gradual adoption of other dermatologic procedures. Lasers for skin rejuvenation were at first ablative (CO₂), but when the long-term complication rates were published, alternatives to this procedure were sought. Nonablative and minimally ablative technologies arose from this search and, in 2006, they are among the most popular procedures performed by dermatologists and those that pretend to be dermatologists alike. Dr. Vic Narurkar will present an update of lasers and light sources.⁵

Changing demographics and improved alternative have increased the numbers of patients with Type IV, V, and VI skin types presenting to dermatologists requesting cosmetic procedures. The advances in lasers, light sources, Botulinum toxins and soft tissue augmentation products directly impact these skin types. In addition to these advances, improved cosmeceuticals now enable physicians to address the unique needs of these skin types. Dr. Jeanine Downie will address these issues.⁶

Cosmeceuticals were once limited to a few vitamins and acids, which had not materially changed in a long time. Recently, however, the advent of potent antioxidants, liposomal-delivery systems, plant-derived growth factors, and a host of other molecules is expanding what is possible with topical esthetic treatments.

Kenneth R. Beer, MD
Guest Editor

References

1. Flynn TC: Update on botulinum toxin. *Semin Cutaneous Med Surg* 25:115-121, 2006
2. Lupo MP: Hyaluronic acid fillers in facial rejuvenation. *Semin Cutaneous Med Surg* 25:122-126, 2006
3. Berlin A, Cohen JL, Goldberg DJ: Calcium hydroxylapatite for facial rejuvenation. *Semin Cutaneous Med Surg* 25:132-137, 2006
4. Beer KR, Rendon MI: Use of Sculptra in esthetic rejuvenation. *Semin Cutaneous Med Surg* 25:127-131, 2006
5. Narurkar VA: Lasers, light sources, and radiofrequency devices for skin. *Semin Cutaneous Med Surg* 25:145-150, 2006
6. Downie JB: Esthetic considerations for ethnic skin. *Semin Cutaneous Med Surg* 25:158-162, 2006